



(11) Publication number:

**0 288 841  
A3**

(12)

**EUROPEAN PATENT APPLICATION**

(21) Application number: **88106017.2**

(51) Int. Cl.<sup>4</sup>: **C07K 3/02 , A61K 37/64**

(22) Date of filing: **15.04.88**

(30) Priority: **27.04.87 US 42925**

(43) Date of publication of application:  
**02.11.88 Bulletin 88/44**

(84) Designated Contracting States:  
**AT BE CH DE ES FR GB GR IT LI NL SE**

(88) Date of deferred publication of the search report:  
**17.05.89 Bulletin 89/20**

(71) Applicant: **MILES INC.**  
**1127 Myrtle Street**  
**Elkhart Indiana 46514(US)**

(72) Inventor: **Shearer, Michael A.**  
**4927 Silverado Drive**  
**Suisun California 94585(US)**

(74) Representative: **Adrian, Albert, Dr. et al**  
**c/o BAYER AG Konzernverwaltung RP**  
**Patentabteilung**  
**D-5090 Leverkusen 1, Bayerwerk(DE)**

(54) **Method of preparing highly purified alpha-1-proteinase inhibitor.**

(57) The invention is related to a highly purified concentrate of alpha-1-proteinase inhibitor and to a method of removing proteins having disulfide bonds, with mild reducing agents and a temperature sufficient to prohibit regrouping of the reduced proteins. This method can be combined with a heat inactivation process.

**EP 0 288 841 A3**



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number

EP 88 10 6017

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
X,D	EP-A-0 067 293 (MEDICAL RESEARCH INSTITUTE OF SAN FRANCISCO) * Page 3, lines 1-37; page 4, lines 1-37; page 5, lines 1-20 *	1-10,12	C 07 K 3/02 A 61 K 37/64
Y,D	---	11	
Y,D	EP-A-0 097 274 (MILES LABORATORIES, INC.) * Page 12, lines 16-31; page 13, lines 1-5 * & US-A-4 379 087	11	
A	ANALYTICAL BIOCHEMISTRY, vol. 124, 1982, pages 364-371, Academic Press, Inc.; C.B. GLASER et al.: "The isolation of alpha-1-protease inhibitor by a unique procedure designed for industrial application" * Front page, abstract *	1-12	
			TECHNICAL FIELDS SEARCHED (Int. Cl.4)
			A 61 K
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 26-02-1989	Examiner FERNANDEZ Y BRANAS F.J.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ----- & : member of the same patent family, corresponding document	

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

(11) Publication number:

**0 288 841  
A2**

(12)

## EUROPEAN PATENT APPLICATION

(21) Application number: 88106017.2

(51) Int. Cl.4: **C07K 3/02 , A61K 37/64**

(22) Date of filing: 15.04.88

(30) Priority: 27.04.87 US 42925

(43) Date of publication of application:  
02.11.88 Bulletin 88/44

(84) Designated Contracting States:  
**AT BE CH DE ES FR GB GR IT LI NL SE**

(71) Applicant: **MILES INC.**  
1127 Myrtle Street  
Elkhart Indiana 46514(US)

(72) Inventor: **Shearer, Michael A.**  
4927 Silverado Drive  
Suisun California 94585(US)

(74) Representative: **Adrian, Albert, Dr. et al**  
c/o **BAYER AG Konzernverwaltung RP**  
Patentabteilung  
D-5090 Leverkusen 1, Bayerwerk(DE)

(54) **Method of preparing highly purified alpha-1-proteinase inhibitor.**

(57) The invention is related to a highly purified concentrate of alpha-1-proteinase inhibitor and to a method of removing proteins having disulfide bonds, with mild reducing agents and a temperature sufficient to prohibit regrouping of the reduced proteins. This method can be combined with a heat inactivation process.

**EP 0 288 841 A2**

## METHOD OF PREPARING HIGHLY PURIFIED ALPHA-1-PROTEINASE INHIBITOR

This invention relates to and has among its objects a method of preparing a high purity alpha-1-proteinase inhibitor (PI) concentrate from blood plasma, blood plasma fractions and concentrates. Further objects of the invention will be evident from the following description wherein parts and percentages are by weight unless otherwise specified.

Alpha-1-proteinase inhibitor is a glycoprotein having molecular weight of 54,000. The protein consists of a single polypeptide chain to which several oligosaccharide units are covalently bound. Human PI has a role in controlling tissue destruction by endogenous serine proteinases. A genetic deficiency of PI, which accounts for 90 % of the trypsin inhibitory capacity in blood plasma, has been shown to be associated with premature development of pulmonary emphysema. The degradation of elastin associated with emphysema probably results from a local imbalance of elastolytic enzymes and the naturally occurring tissue and plasma proteinase inhibitors. PI rapidly inhibits human pancreatic and leukocyte elastases (Biochem. Biophys. Res. Comm., Vol. 72, No. 1, pages 33-39, 1976; ibid., Vol. 88, No. 2, pages 346-350, 1979).

A number of methods employed to isolate PI from blood plasma are set forth in U.S. Patents 4,439,358, 4,379,087, and 4,656,254 and are incorporated herein by reference.

Pannell et al., Biochemistry, Vol. 13, pages 5439-5445, (1974), teaches a method of depleting albumin from plasma using Sepharose-Blue Dextran adsorption are then pooled and fractionated with solid ammonium sulfate.

Glaser et al., EP 82103473, teach the use of reducing agents, such as dithiothreitol to break disulfide bonds and precipitation by salting out with 50%  $(\text{NH}_4)_2\text{SO}_4$ , to remove unwanted proteins.

The invention described herein is a method of preparing a highly purified PI concentrate by depleting the albumin from a solution containing PI by adding a mild reducing agent sufficient to break the disulfide bonds in the albumin and heating the solution containing the PI and mild reducing agent to prevent the destabilized proteins from recombining.

In preferred embodiments, PI is separated from blood Plasma or blood plasma fractions according to the Coan methods, U.S. Patents 4,379,087 and 4,439,358 and U.S. Serial No. 793,807. As well as the improvements set forth in U.S. Patent 4,656,254, Improved Methods of Preparing alpha-1-proteinase inhibitor. The reducing agent can be added to a solution of separated PI to break any protein disulfide bonds without reducing the PI, since PI does not have disulfide bonds. Reducing

agents which can be utilized are dithiothreitol, dithioerythritol, 2-mercaptoethanol, mercaptoacetic acid, sodium borohydride and cysteine. The reducing agents are present in a concentration of about 2mM to 8mM. The concentration of reducing agent will vary depending on the type of solution containing PI and the choice of reducing agent. The amount of reducing agent should be sufficient to reduce the disulfide bonds without denaturing or damaging PI or any other protein to be recovered. The pH of the solution containing PI and the reducing agent can be adjusted to optimize the rate of reaction and also to stop the reaction. Heat denaturation of the reduced protein can be accomplished by raising the temperature to about 35°C or higher.

Suitable pH ranges are of about 7.9 to about 8.5 whereas the rates of reaction are about 5 to about 95 minutes.

Heat denaturation can also be accomplished in combination with a heat treating viral inactivation step. In U.S. Patent 4,470,968, there is disclosed a method of stabilizing coagulation Factors II, VIII, IX, and X against heat during pasteurization or heating at a temperature of about 60°-100°C by mixing with heat-stabilizing or pasteurization-stabilizing, amounts of a polyol and a source of citrate ions. Pasteurized compositions of these coagulation Factors were obtained by heating a mixture of unpasteurized protein composition, a polyol, and a source of citrate ions suspended or solubilized usually in an aqueous medium at a temperature and for a time sufficient to pasteurize the protein composition. Following pasteurization or heat treatment, the polyol and citrate ions are removed totally, or in part, as desired, from the protein composition by conventional techniques, and the pasteurized protein composition is processed according to conventional procedures for its ultimate therapeutic use. By way of illustration and not limitation, one may employ the method of U.S. Patent No. 4,470,968 (herein incorporated by reference) or any of the methods of the prior art described therein for heat inactivation.

One advantage of the invention is the availability of an essentially pure PI preparation that does not require the addition of ammonium sulfate to remove unwanted protein.

Another advantage of the invention is the incorporation of a step to remove unwanted protein with a heat inactivation step. The mild reducing agent can be added to the aqueous solution any time prior to the heat inactivation step and unwanted proteins removed when the pasteurized protein composition is processed according to convention-

al procedures for ultimate therapeutic use.

The reducing agent and denatured proteins can be removed by ultrafiltration and anion exchange methods without the addition of a salting out step.

Current methods for obtaining PI from blood plasma use as a starting material: Cohn Fraction IV and IV-1 and includes reworks of these fractions wherein other proteins are first removed, Cohn Effluent II + III and Cohn Effluent I and cryosupernatant solution. Blood plasma is fractionated according to the Cohn ethanol fractionation technique or its modifications [see for example, Cohn *et al.*, *J. Am. Chem. Soc.*, 68,459, (1946); Oncley *et al.*, *ibid.*, 71,541 (1949); U.S. Patent 2,390,074; "The Plasma Proteins", second edition, Vol. III, pages 548-550, Academic Press, New York, N.Y.] to give a Cohn fraction containing PI. This method for removing proteins containing disulfide bonds could have application in genetically engineered PI, in those situations where the solution containing PI, such a tissue culture medium or other eluates, also contained disulfide proteins, such as albumin.

PI was produced following the method outlined by Coan *et al.*, *Vox Sang.* 48:333-342, 1985; U.S. Patents 4,379,087 and 4,439,358, and improvements set forth by Shearer *et al.*, U.S. Patent 4,656,254.

Dithiothreitol was added to Coan's purified column eluate at a concentration of 5mM, pH was adjusted to 8.0-8.2. The solution was allowed to mix for twenty minutes and the reaction was stopped by adjusting the pH to 6.5. The PI solution containing the mild reducing agent was heat treated at 60°C for 10 hours in the presence of sucrose and citrate as indicated by Coan and Mitra, *Vox Sang.* 46:142-148 (1984) to render the solution free of virus. The denatured proteins were removed by applying the heat-treated solution to a DEAE Sepharose column equilibrated with 0.025M sodium phosphate at pH 6.5. The PI was removed by using a phosphate gradient to 0.1M at pH 6.5. The PI was found to be approximately 100% pure by Specific Activity (mg PI/A<sub>280</sub>), CAE and HPLC. In contrast, the prior art method of Coan/Shearer provided a PI purity of about 60%.

The invention is related to a method of preparing highly purified alpha-1-proteinase inhibitor from a mixture of blood plasma proteins comprising

(a) adding a mild reducing agent to a mixture of plasma proteins containing alpha-1-proteinase inhibitor, said reducing agent being in a concentration sufficient to reduce disulfide bonds without reducing alpha-1-proteinase inhibitor;

(b) raising the temperature of the solution of step (a) to above 35°C; and

c) recovering alpha-1-proteinase inhibitor from the solution.

The above method of removing unwanted pro-

teins from a mixture of plasma proteins containing alpha-1-proteinase inhibitor can be accomplished in modified form by the steps comprising

(a) adding a mild reducing agent to a mixture of plasma proteins containing alpha-1-proteinase inhibitor, said reducing agent being in a concentration sufficient to reduce disulfide bonds without reducing alpha-1-proteinase inhibitor;

(b) mixing the composition of Step (a) with a polyol selected from sucrose or reduced sugars and a source of citrate ions in an aqueous medium, said polyol and citrate ions being present in an amount sufficient to stabilize the protein during pasteurization;

(c) and heating the mixture at a temperature and for a time sufficient to pasteurize the composition.

Generally the amount of polyol and citrate ions in the mixture of Step (b) is about 20% to saturation of polyol. On a weight to volume basis and about 0.1-1.0 moles of citrate ion per liter of solution.

Heat denaturation can be accomplished independently of a viral inactivation step by raising the temperature of the solution containing PI and the mild reducing agent sufficient to reduce the disulfide bonds to a level above 35°C. The reducing agent can be removed by ultrafiltration. PI can be recovered from the heat treated solution by column chromatography using an appropriate anion exchange medium.

The mixture of the latter mentioned method is heated in step (c) at a temperature of about 60° - 100°C for a period of about 1 - 10 hours. The method further includes the Step of removing the polyol and citrate ions from the mixture of Step (c), for instance, by subjecting the mixture to diafiltration or by subjecting the mixture to dialysis or ion exchange chromatography. The concentration of citrate ions is about 0.3M to about 0.5M. A preferred embodiment is where the concentration of sucrose is, for instance, 37% (w/v) and the concentration of citrate ions is 0.38M.

The aqueous solution of plasma proteins containing alpha-1-proteinase inhibitor is preferably selected from the group of plasma fractions consisting of Cohn Fraction IV-1, Cohn Fraction IV, reworks of Cohn Fraction IV and IV-1, Cohn Effluent II + III, Cohn Effluent I, and cryosupernatant solution.

Especially the aqueous solution of plasma proteins containing alpha-1-proteinase inhibitor is selected from the group consisting of Cohn Fraction IV-1, Cohn Fraction IV, Cohn Effluent II + III and Cohn Effluent I.

Most preferred is a method wherein the aqueous solution of plasma proteins containing alpha-1-proteinase inhibitor is Cohn Fraction IV-1.

The invention is also related to a method for separating alpha-1-proteinase inhibitor from an aqueous solution of plasma proteins containing alpha-1-proteinase inhibitor which comprises the steps of:

(a) contacting an aqueous solution of one of the group of blood plasma and a blood fraction containing alpha-1-proteinase inhibitor with from about 8% to about 23% (w/v), based on volume of the aqueous solution, of a polycondensed polyalkylene glycol selected from polyethylene glycol and polypropylene glycol, at a temperature of from about 2°C to about 50°C for a period of about 0.2 - 24 hours to selectively precipitate unwanted proteins from the solution without precipitating alpha-1-proteinase inhibitor to obtain a mixture containing alpha-1-proteinase inhibitor free of unwanted proteins,

(b) recovering the mixture from step (a), and

(c) separating alpha-1-proteinase inhibitor from the mixture recovered in step (b), by column chromatography providing an eluate containing alpha-1-proteinase inhibitor, which is characterised by the step of:

(i) adding a mild reducing agent to a mixture of plasma proteins containing alpha-1-proteinase inhibitor, said reducing agent being in a concentration sufficient to reduce disulfide bonds without reducing alpha-1-proteinase inhibitor;

(ii) raising the temperature of the solution of step (a) to above 35°C; and

(iii) recovering alpha-1-proteinase inhibitor from the solution.

The method according to the process at page 5 further includes the steps of

(i) providing for use in step (a) an aqueous solution of the one of the blood plasma and blood plasma fraction dissolved in from 20 to 100 volumes of buffer-solution per weight of the plasma or plasma fraction used, and

(ii) prior to use in step (a), adjusting one of the buffer solution and the resulting aqueous solution of the blood plasma and blood plasma fraction from step (1) to render the pH of the resulting solution at from 9.0 to 11.0.

Preferably in step (i) there is used from 20 to 50 volumes of buffer solution per weight of the one of the plasma and plasma fraction.

More preferred in step (i) there is used from 20 to 30 volumes of buffer solution per weight of the one of the plasma and plasma fraction and most preferred in step (i) there is used 24 volumes of buffer solution per weight of the one of the plasma and plasma fraction.

The pH of the one of the buffer solution and the resulting aqueous solution is adjusted to render the pH of the resulting solution from 9.0 to 10.5.

Preferably the pH of the one of the buffer

solution and the resulting aqueous solution is adjusted to render the pH of the resulting solution from 9.0 to 10.5 and, more preferred the pH of the one of the buffer solution and the resulting aqueous solution is adjusted to render the pH of the resulting solution from 9.0 to 10.5.

## Claims

1. A method of preparing highly purified alpha-1-proteinase inhibitor from a mixture of blood plasma proteins comprising

(a) adding a mild reducing agent to a mixture of plasma proteins containing alpha-1-proteinase inhibitor, said reducing agent being in a concentration sufficient to reduce disulfide bonds without reducing alpha-1-proteinase inhibitor;

(b) raising the temperature of the solution of step (a) to above 35°C; and

(c) recovering alpha-1-proteinase inhibitor from the solution.

2. The method of claim 1 wherein said mild reducing agent is selected from the group consisting of dithiothreitol, dithioerythritol, sodium borohydride, beta-mercaptoethanol, and cysteine.

3. The method of claim 2 wherein the concentration of said reducing agent is in the range of about 2mM to about 8mM.

4. The method of claim 3 wherein said reducing agent is 5mM dithiothreitol.

5. The method of claim 2 wherein said solution is mixed from about 5 to about 95 minutes at a pH range of about 7.9 to about 8.5.

6. The method of claim 5 wherein the temperature range is above 35°C.

7. The method of claim 4 wherein said solution is mixed for a period of about 20 minutes at a pH range of about 8.0 to about 8.2 at a temperature of about 25°C.

8. The method of claim 2 wherein said reaction is stopped by adjusting the pH to below 7.8.

9. The method of claim 8 wherein the pH is adjusted to about 6.5.

10. A method according to claim 1 including the further step of treating said aqueous solution of alpha-1-proteinase inhibitor obtained from step (c) to inactivate, and to reduce the infectivity of, infectious microorganisms so as to render the alpha-1-proteinase inhibitor non-infectious to such infectious microorganisms and thereby render the alpha-1-proteinase inhibitor useful for therapeutic and prophylactic purposes.

11. A method of preparing high purified alpha-1-proteinase inhibitor from a mixture of blood plasma proteins comprising

(a) adding a mild reducing agent to a mixture of plasma proteins containing alpha-1-proteinase inhibitor, said reducing agent being in a concentration sufficient to reduce disulfide bonds without reducing alpha-1-proteinase inhibitor;

5

(b) mixing the composition of Step (a) with a polyol selected from sucrose or reduced sugars and a source of citrate ions in an aqueous medium, said polyol and citrate ions being present in an amount sufficient to stabilize the protein during pasteurization;

10

(c) and heating the mixture at a temperature and for a time sufficient to pasteurize the composition.

12. A pharmaceutical composition comprising a protease inhibitory effective amount of alpha-1-proteinase inhibitor produced by the method according to claims 1 to 11 and a pharmaceutically acceptable carrier.

15

20

25

30

35

40

45

50

55

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

(11) Publication number:

**0 288 841**  
**A3**

(12)

## EUROPEAN PATENT APPLICATION

(21) Application number: 88106017.2

(51) Int. Cl.4: C07K 3/02 , A61K 37/64

(22) Date of filing: 15.04.88

(30) Priority: 27.04.87 US 42925

(43) Date of publication of application:  
02.11.88 Bulletin 88/44(84) Designated Contracting States:  
AT BE CH DE ES FR GB GR IT LI NL SE(88) Date of deferred publication of the search report:  
17.05.89 Bulletin 89/20(71) Applicant: MILES INC.  
1127 Myrtle Street  
Elkhart Indiana 46514(US)(72) Inventor: Shearer, Michael A.  
4927 Silverado Drive  
Suisun California 94585(US)(74) Representative: Adrian, Albert, Dr. et al  
c/o BAYER AG Konzernverwaltung RP  
Patentabteilung  
D-5090 Leverkusen 1, Bayerwerk(DE)

(54) Method of preparing highly purified alpha-1-proteinase inhibitor.

(57) The invention is related to a highly purified concentrate of alpha-1-proteinase inhibitor and to a method of removing proteins having disulfide bonds, with mild reducing agents and a temperature sufficient to prohibit regrouping of the reduced proteins. This method can be combined with a heat inactivation process.

**EP 0 288 841 A3**





European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number

EP 88 10 6017

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
X,D	EP-A-0 067 293 (MEDICAL RESEARCH INSTITUTE OF SAN FRANCISCO) * Page 3, lines 1-37; page 4, lines 1-37; page 5, lines 1-20 *	1-10, 12	C 07 K 3/02 A 61 K 37/64
Y,D	---	11	
Y,D	EP-A-0 097 274 (MILES LABORATORIES, INC.) * Page 12, lines 16-31; page 13, lines 1-5 * & US-A-4 379 087	11	
A	ANALYTICAL BIOCHEMISTRY, vol. 124, 1982, pages 364-371, Academic Press, Inc.; C.B. GLASER et al.: "The isolation of alpha-1-protease inhibitor by a unique procedure designed for industrial application" * Front page, abstract *	1-12	
			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
			A 61 K
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 26-02-1989	Examiner FERNANDEZ Y BRANAS F.J.
<b>CATEGORY OF CITED DOCUMENTS</b> X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ----- & : member of the same patent family, corresponding document			